

TYROSINE HYDROXYLASE IMMUNOREACTIVE FIBERS IN THE UPPER LAYERS OF
THE ADOLESCENT RAT MEDIAL PREFRONTAL CORTEX

BY

JOSEPH MICHAEL BRODSKY

THESIS

Submitted in partial fulfillment of the requirements
for the degree of Master of Science in Biology
in the Graduate College of the
University of Illinois at Urbana-Champaign, 2015

Urbana, Illinois

Adviser:

Professor Janice Juraska

Abstract

Adolescence is an important period of maturation for the brain. In particular, the medial prefrontal cortex continues developing throughout adolescence and is responsible for a number of executive and cognitive functions. Innervation of the medial prefrontal cortex by the dopaminergic system contributes to some of these maturing behaviors and the dopaminergic system is implicated in mental illnesses. However, much previous work on medial prefrontal cortex development over adolescence has considered only one sex or has failed to capture timing of changes. Here, the development of tyrosine hydroxylase-immunoreactive axons, to indicate dopaminergic innervation, in the upper layers of the medial prefrontal cortex was examined in both sexes of Long-Evans rats during five ages from postnatal day 25 to 90. Tyrosine hydroxylase-immunoreactive axon density was found to increase over the course of adolescence, with the most marked increases being prepubertal in both males and females. No sex differences were identified. These results suggest a similar development, and thus similar susceptibility to disruption, of the dopaminergic system in the rat medial prefrontal cortex in both sexes.

Table of Contents

Introduction.....	1
Methods.....	4
Animals.....	4
Tissue Preparation.....	4
Immunocytochemistry.....	4
TH Analysis.....	5
Statistical Analysis.....	6
Results.....	7
Layer 1.....	7
Layer 2/3.....	10
Discussion.....	14
Acknowledgments.....	18
List of References.....	19

Introduction

Adolescence is an important period of brain development for humans and other mammals. During this time, neuroanatomical changes include pruning and refinement of neurons, dendrites, synapses, and expression of receptors (Markham et al., 2007; Lewis, 1997; Huttenlocher and Dabholkar, 1997; Andersen et al., 2000). Changes during this time can show sex differences and may or may not involve sex hormones (Juraska et al., 2013). In humans, anatomical changes produce an observable volume change using MRI, with a sex difference in age of peak volume (Giedd et al., 1999). Behavioral changes are also pronounced, with changes in many executive and social behaviors, notably including decisions involving risk-taking and reward (Spear, 2000). Adolescence itself is difficult to delimit, but in humans it occurs in the second decade of life, and is often considered to be from 12 to 18 years of age, or even older; in rats, postnatal day (P)28 to P42 or P45 can be considered adolescence, although some changes continue to P60, such as increasing connections from the amygdala to the prefrontal cortex (Spear, 2000; Brenhouse and Andersen, 2011; Cunningham et al., 2002).

The prefrontal cortex, particularly the medial prefrontal cortex (mPFC), is an important brain region for executive behavior (Dalley et al., 2004), including decision making based on reward and cognitive flexibility. Many of these prefrontal behaviors mature during adolescence, with adolescents displaying increased impulsivity and lower cognitive flexibility than adults (Steinberg, 2010; Andrzejewski et al., 2011). Anatomically, the prefrontal cortex undergoes changes including cell number, synaptic density, connectivity to other brain regions, and receptor expression over the course of adolescence in both rats and humans (Markham et al., 2007;

Huttenlocher and Dabholkar, 1997; Cunningham et al., 2002; Andersen et al., 2000).

Dopaminergic innervation to the mPFC contributes to the functions of the mPFC. Tyrosine hydroxylase (TH) is the rate-limiting step in the synthesis of dopamine, an important catecholamine neurotransmitter, and is commonly used to mark dopaminergic fibers (e.g. Lewis et al., 1987; Naneix et al., 2012). Dopaminergic axons from the ventral tegmental area target the mPFC in rodents and primates, and this innervation increases over adolescence (Kalsbeek et al., 1988; Lewis et al., 1987; Naneix et al., 2012). Furthermore, dopamine receptor expression peaks during late adolescence (Andersen et al., 2000). The response of neurons to dopamine also changes during this time (Tseng and O'Donnell, 2007). Dopamine in the mPFC plays a role in adult behaviors that matured over adolescence, such as cognitive flexibility (Gruber et al., 2010).

Mental illness is also associated with adolescence. The highest rates of onset of many psychopathologies, including mood disorders and schizophrenia, occur during this time (Kessler et al., 2005). Additionally, there are sex differences in disease onset, with more adolescent males experiencing schizophrenia and females experiencing depression (Abel et al., 2010; Parker and Brotchie, 2010). Exposure to stress can expose or exacerbate these disorders (Arnsten, 2011; Monroe and Harkness, 2005). These illnesses involve the prefrontal cortex. Schizophrenia shows reductions in prefrontal cortex volume and synapses (Lewis, 1997; Giedd et al., 2008); depression also involves changes in prefrontal cortex volume and changes in activity (Palazidou, 2012). Furthermore, the dopamine system is implicated in both depression and schizophrenia (Dunlop and Nemeroff, 2007; Money and Stanwood, 2013). Combined, these data suggest that disruption of normal development of prefrontal cortex may produce or exacerbate psychopathologies (Giedd et al., 2008).

Understanding normal adolescent development is important for an appreciation of how the adult brain forms and for disease models. However, studies of adolescent development often use only one sex (e.g. Andersen et al., 2000), few timepoints (e.g. Markham et al., 2007), or methods that reveal little about the bases for observed change (e.g. Giedd et al., 1999). Previous studies of dopaminergic innervation in the mPFC have been no exception and in particular relied on male animals only (e.g. Naneix et al., 2012). The following work aims to elucidate some of the changes occurring during this important period. Here, we examine the development of TH-immunoreactive fibers in the upper layers of the adolescent rat mPFC at several ages, to identify age and sex differences from preadolescence to adulthood. These timepoints intend to capture changes over adolescence, with P25 being preadolescent and P90 adult. Female rats enter puberty before male rats, with puberty occurring by P35 in females, with males undergoing puberty after P35 but typically finishing by P45. Since development can continue after puberty, P60 is also examined as late adolescence.

Methods

Animals

Long-Evans hooded rats (Harlan) were bred in the Psychology Department's vivarium. Following weaning at P24, same-sex littermates were group housed (2-3) in transparent plastic cages in a temperature-controlled room with 12:12h light-dark and with ad lib. access to a standard rat chow and water. Animals were checked for puberty markers between P25-48. Animal experimentation was approved by the University of Illinois' Institutional Animal Care and Use Committee.

Tissue Preparation

At P25, P35, P45, P60, or P90, 10-11 rats of each sex (total: 102 animals) were anesthetized with sodium pentobarbital (100mg/kg) and transcardially perfused with 0.1M phosphate-buffered saline (PBS) followed by 4% paraformaldehyde in PBS. Brains were removed and stored in paraformaldehyde solution for 24 hours, then transferred to a 30% sucrose solution. After 3 days, brains were cut on a freezing microtome into 40µm-thick coronal sections. Every fifth section was used for TH immunocytochemistry; other sections were Nissl stained or saved for further experiments. Sections were stored in 30% glycerol, 30% ethylene glycol, 10% PBS, and dH₂O.

Immunocytochemistry

For TH labelling, sections were rinsed in tris-buffered saline (TBS; pH 7.6), placed in a

blocking solution (1% hydrogen peroxide, 20% normal goat serum, 1% BSA in TBS) for 30 minutes, and then exposed to monoclonal mouse anti-tyrosine hydroxylase (Millipore) at 1:1000 dilution in TTG (2% normal goat serum and 0.3% triton-x-100 in TBS) for 48 hours followed by a rinse in TTG, antimouse biotinylated antibodies (Vector Labs) in TTG for 90 minutes, rinses in TTG and TBS, avidin-biotin-complexed horseradish peroxidase (Vector Labs) for 1 hour, rinsed in TBS, diaminobenzidine (Sigma Fast Tabs), and finally rinsed in TBS, and then mounted on slides, allowed to dry, and then coverslipped.

TH Analysis

To quantify TH fibers, a Zeiss Axiovert 200M (Carl Zeiss, Thornwood, NY) with a 63x oil immersion objective was used to obtain Z-stacks of layers 1, 2/3, and 5/6 from both left and right hemispheres in two sections per brain within the mPFC. The mPFC was defined as containing the prelimbic and infralimbic regions, starting rostrally with the section where white matter first occurs and ending when the genu of the corpus callosum is encountered (Chisholm et al., 2012; for cytoarchitectonic parcellation of the mPFC, Markham et al., 2007). Images in the stacks were taken every 0.275 μ m, resulting in around 100 images per stack. Stacks were taken approximately midway between the dorsal and ventral extent of white matter, while avoiding obvious artifacts. Layer 1 was taken as close to the medial surface of the mPFC as possible (the images contained the entire layer), layers 2/3 close to layer 1 while excluding layer 1 fibers, and layers 5/6 in the middle of 5/6. The layers have readily identifiable arrangements of fibers: layer 1 contains distinctive dorsoventally-aligned fibers, layers 2/3 contain some prominent lateral fibers, and layers 5/6 contain numerous fibers running in all directions. Z-stacks were refocused to a single image using real wavelets with the Extended Depth of Field plugin (Forster et al.,

2004) for ImageJ/FIJI (Schneider et al., 2012; Schindelin et al., 2012). Processed images were then examined for extraneous staining, which were excluded by hand with a white brush in ImageJ. Items removed were blood vessels, with membranous, weakly-staining, hollow, or segmented appearance; diffuse background staining; large amorphous or soma-like objects; dirt and tissue artifacts, which were grey, black, or blue rather than brown; and blobs or putative cells which did not appear collinear with presumptive fibers. The images were converted to binary images with black for stained fibers and the area covered in black pixels was quantified before and after processing. The binary processed image was then skeletonized, reducing contiguous black pixels to structures 1 pixel wide, and this area was also quantified. Skeletonizing the image preserves fiber length while unskeletonized images quantify differences in fiber thickness and length.

Because the mPFC changes in size across the ages examined, the percent of image stained was also multiplied by the volume of the appropriate mPFC layers to produce μm^3 of stained material. The volumes were obtained from Nissl stained sections parcellated stereologically by other experimenters as in Markham et al. (2007).

This work deals with layers 1 and 2/3; another experimenter is quantifying layers 5/6.

Statistical Analysis

Statistical analysis was performed using SPSS (IBM). ANOVAs were performed using age and sex as factors, with litter and cohort as covariates. Preplanned post hoc comparisons (t-tests) were examined between ages P25-35, P35-45, P45-60, and P60-90 when age was significant.

Results

Layer 1

The ANOVA of processed binary images (density: number of TH labeled pixels/volume) (Figure 1) revealed an effect of age ($F(4, 90)=5.630$; $p<.001$) and immunohistochemistry cohort ($F(1, 90)=34.619$; $p<.001$), but not sex ($F(1, 90)=.366$; $p=.547$) nor an age by sex interaction ($F(4, 90)=1.024$; $p=.400$). Likewise, skeletonized images (Figure 2) showed the same age ($F(4, 90)=3.951$; $p=.005$) but not sex ($F(1, 90)=.259$; $p=.612$) effect, with no interaction ($F(4, 90)=.488$; $p=.744$). The same pattern of effects was present in the unprocessed images (age: $F(4, 90)=4.949$; $p=.001$; sex: $F(1, 90)=1.111$; $p=.295$; age by sex: $F(4, 90)=.730$; $p=.574$) (Figure 3).

When percent of image stained was multiplied by volume of layer 1 to produce the amount (μm^3) of stained axons, a main effect of age persisted for processed images in males ($F(4, 44)=3.375$; $p=.017$) and females ($F(4, 44)=2.723$; $p=.041$) (Figure 4). Post hoc comparisons of the amount of stained axons showed a significant difference between P25 and P35 in females ($p=.030$) and a trend in males ($p=.060$); all other post hocs were not significant. Only trends in skeletonized images for males ($F(4, 44)=2.390$; $p=.065$) and females ($F(4, 44)=2.092$; $p=.098$) were found (Figure 5).




Figure 1: Percent stained of binary processed layer 1 images. Significant effect of age: $F(4, 90)=5.630$; $p<.001$.

Means + SEM.




Figure 2: Percent stained of skeletonized layer 1 images. Significant effect of age: $F(4, 90)=3.951$; $p=.005$. Means

+ SEM.




Figure 3: Percent stained of unprocessed layer 1 images. Significant effect of age: $F(4, 90)=4.949$; $p=.001$. Means + SEM.




Figure 4: Volume (μm^3) of stained axons from binary processed layer 1 images. Effect of age in males: $F(4, 44)=3.375$, $p=.017$; and females: $F(4, 44)=2.723$, $p=.041$. *: $p<.05$; #: $p<.10$. Means + SEM.




Figure 5: Volume (μm^3) of stained axons from layer 1 skeletonized images. Trend of age for males: $F(4, 44)=2.390$, $p=.065$; and females: $F(4, 44)=2.092$, $p=.098$. Means + SEM.

Layer 2/3

There was an effect of age ($F(4, 90)=3.103$; $p=.019$) in the processed binary images of density (Figure 6) and an effect of immunohistochemistry cohort ($F(1, 90)=13.484$; $p<.001$) but not sex ($F(1, 90)=.315$; $p=.576$) nor age by sex ($F(4, 90)=.683$; $p=.606$). Similarly, skeletonized images (Figure 7) showed age ($F(4, 90)=3.237$; $p=.016$) but not sex ($F(1, 90)=.045$; $p=.832$) nor age by sex ($F(4, 90)=.547$; $p=.702$). Unprocessed images (Figure 8) showed similar results (age: $F(4, 90)=2.596$; $p=.042$; sex: $F(1, 90)=.477$; $p=.492$; age by sex: $F(4, 90)=.579$; $p=.679$).

When the volume (μm^3) of stained axons was considered, binary processed images (Figure 9) showed an effect of age in males ($F(4, 44)=2.939$; $p=.031$) and females ($F(4, 42)=2.890$; $p=.034$). Post hoc tests again showed a significant difference only between P25 and P35 in males ($p=.006$) and only a trend in binary images in females between P25 to P35

($p=.062$). Skeletonized (Figure 10) also showed an effect of age in males ($F(4, 44)=3.268$; $p=.020$) but a trend in females ($F(4, 42)=2.292$; $p=.075$). Post hocs revealed a significant difference between P25 and P35 in males ($p=.002$).




Figure 6: Percent stained of binary processed layer 2/3 images. Significant effect of age: $F(4, 90)=3.103$; $p=.019$.

Means + SEM.




Figure 7: Percent stained of skeletonized layer 2/3 images. Significant effect of age: $F(4, 90)=3.237$; $p=.016$.

Means + SEM.




Figure 8: Percent stained of unprocessed layer 2/3 images. Significant effect of age: $F(4, 90)=2.596$; $p=.042$.

Means + SEM.




Figure 9: Volume (μm^3) of stained axons from binary processed layer 2/3 images. Significant effect of age in males:

$F(4, 44)=2.939$, $p=.031$; and females: $F(4, 42)=2.890$, $p=.034$. *: $p<.05$; #: $p<.10$. Means + SEM.




Figure 10: Volume (μm^3) of stained axons from skeletonized layer 2/3 images. Effect of age in males: $F(4,$

$44)=3.268$; $p=.020$. Trend of age in females: $F(4, 42)=2.292$, $p=.075$. *: $p<.05$. Means + SEM.

Discussion

The work presented here finds no significant sex differences in the increase of TH immunoreactive axons in the rat mPFC, and with much of the increase seeming to occur prepubertally in both sexes.

Previous work has shown an increase in TH density, and in the dopaminergic system, in the mPFC over the course of adolescence in male rats in particular. Kalsbeek et al. (2008) qualitatively described mPFC dopamine-immunoreactive fibers from embryonic development to P60 and P90, but did not report potential sex differences from their mixed-sex subjects. They found an increase in dopamine-containing axons between P20 and P60, but no changes from P60 to P90, and they report the increase as being obvious in the upper layers of prelimbic mPFC. Naneix et al. (2012) examined a number of dopaminergic features in the mPFC, but used only male rats. They found a gradual increase in TH-immunoreactive fibers in the mPFC from P25 to P70, along with an increase in cortical dopamine between P45 and P70 and a peak in the expression of several dopamine receptor mRNAs around P45. Other studies report overproduction of dopamine receptors in late adolescence in the male rat mPFC (Andersen et al., 2000). However, the possibility of sex differences in the development of this system has not been well-studied.

The timing of TH increases observed here suggests that these changes are not mediated by pubertal hormones in male rats. Androgens surge around P45 with the onset of puberty in males. Since no hormonal quantification or manipulation was performed, whether pubertal hormones in females ramping up around P35 occurred early enough to affect TH cannot be ruled

out. That males and females may have different mechanisms underlying development is not unheard of. Changes in neuron number in the rat mPFC over adolescence have been associated with ovarian but not testicular hormones (Koss et al., 2015). Adult animals show effects of gonadal hormones on TH. Adult male rats show increased TH density in prefrontal cortex following orchiectomy which is reduced with testosterone but not estrogen (Kritzer et al., 2007). A decrease in TH density caused by androgens during adolescence might be overwhelmed and hidden by stronger coincident growth of fibers resulting in a net increase in density, or the neurons may respond differently at this stage of development so that androgens do not negatively influence density. In adult rats, some ventral tegmental dopaminergic neurons that project to the mPFC express androgen receptors and a smaller number express estrogen receptor β (Kritzer and Creutz, 2008), and counts of TH positive neurons in the ventral tegmental area can change by P90 following orchiectomy on P60 (Johnson et al., 2010), suggesting androgens can influence the dopaminergic system before the final age examined here. Whether pubertal hormones affect the adolescent development of TH-immunoreactive projections to the mPFC directly, indirectly, or not at all, the dopaminergic system displays sex differences and may respond to perinatal and adult hormonal milieu (Gillies et al., 2014).

The present work considers only the upper layers of the cortex. However, TH immunoreactive fiber density is greater in layer 5/6 in the mPFC (Descarries et al., 1987). It is not improbable that the deeper layers may show more pronounced changes. Position within the layers of the cortex might not be the only influence on degree of observed change. When extremely rostral or caudal sections of the male rat mPFC are considered, an effect of age is seen in rostral, but not caudal, sections over adolescence (Naneix et al., 2012). The present work used

sections from about the same rostrocaudal level in each animal, about midway through the mPFC; the effects seen here may be muted or amplified compared to more extreme sections.

TH is used as a marker for dopaminergic neurons, but other cells may be immunoreactive. Besides background staining, noradrenergic axons project throughout the cortex and may be responsible for much cortical dopamine (Devoto and Flore, 2006). These fibers, however, are generally weakly staining for TH, depending on the antibody used (Verney et al., 1982; Verney et al., 1993; Lewis et al., 1987; Miner et al., 2003; Zhang et al., 2011). The rat mPFC is notable for being richly innervated by dopaminergic fibers but is not especially enriched in noradrenergic fibers compared to the rest of the neocortex (Devoto and Flore, 2006). Other studies which stained for dopamine β -hydroxylase, which converts dopamine to norepinephrine, did not find an increase in mPFC noradrenergic fibers in male rats over adolescence (Naneix et al., 2012), or found distinct populations of TH fibers in primates where TH fibers survived lesioning of noradrenergic pathways (Lewis et al., 1987). Lesions of noradrenergic pathways do not significantly lower TH levels in medial prefrontal cortex (Emson and Koob, 1978). Tritiated norepinephrine reveals adult levels of noradrenergic fiber density in non-mPFC cortex prepubertally (Levitt and Moore, 1979). Thus, it is not thought that incidental staining of noradrenergic fibers has significantly influenced the present study. Additionally, a few interneurons in the cerebral cortex express TH but do not produce catecholamines and lose immunoreactivity with age (Asmus et al., 2008; Asmus et al., 2011). The location, and even presence, of these cells varies by species but in rats they tend to reside in layer 2/3 (Sato and Suzuki, 1990; Benavides-Piccione and DeFelipe, 2007; Asmus et al., 2008) and are most common in a postnatal window from P16 ending around P40 (Asmus et al., 2008). It is not

expected that these cells affect the present results because they peak prepubertally and decrease rapidly.

The development of the TH innervation of the mPFC continues to pose questions. Since the increases in TH-immunoreactive fibers did not show sex differences and had the most significant gains prepubertally, insults to this system may be most disruptive when presented early. Barring hidden sex differences, male and female rats would be expected to display similar disruption to TH innervation and similar deficits in dopamine-related mPFC behaviors. How the development of the mPFC and sex differences in pathological outcomes are related remains to be clarified.

Acknowledgments

The following people contributed to this project: Janice Juraska, Jari Willing, Taehyeon Kim, Laura Cortes, Carly Drzewiecki. Funding was provided by NIH grant MH099625. Imaging was done at the Microscopy Suite in the Beckman Institute.

List of References

Abel KM, Drake R, Goldstein JM (2010) Sex differences in schizophrenia. *Int Rev Psychiatry* 22(5):417-428.

Andersen SL, Thompson AT, Rutstein M, Hostetter JC, Teicher MH (2000) Dopamine receptor pruning in prefrontal cortex during the periadolescent period in rats. *Synapse* 37:167-169.

Andrejewski ME, Schochet TL, Feit EC, Harris R, McKee BL, Kelley AE (2011) A comparison of adult and adolescent rat behavior in operant learning, extinction, and behavioral inhibition paradigms. *Behav Neurosci* 125(1):93-105.

Arnsten AFT (2011) Prefrontal cortical network connections: key site of vulnerability in stress and schizophrenia. *Int J Dev Neurosci* 29(3):215-223.

Asmus SE, Anderson EK, Ball MW, Barnes BA, Bohnen AM, Brown AM, Hartley LJ, Lally MC, Lundblad TM, Martin JB, Moss BD, Phelps KD, Phillips LR, Quilligan CG, Steed PR, Terrell SL, Warner AE (2008) Neurochemical characterization of tyrosine hydroxylase-immunoreactive interneurons in the developing rat cerebral cortex. *Brain Res* 1222:95-105.

Asmus SE, Cocanougher BT, Allen DL, Boone JB, Brooks EA, Hawkins SM, Hench LA, Ijaz T, Mayfield MN (2011) Increasing proportions of tyrosine hydroxylase-immunoreactive interneurons colocalize with choline acetyltransferase or vasoactive intestinal peptide in the developing rat cerebral cortex. *Brain Res* 1383:108-119.

Benavides-Piccione R, DeFelipe J (2007) Distribution of neurons expressing tyrosine hydroxylase in the human cerebral cortex. *J Anat* 211:212-222.

Brenhouse HC, Andersen SL (2011) Developmental trajectories during adolescence in males and females: a cross-species understanding of underlying brain changes. *Neurosci Biobehav Rev* 35(8):1687-1703.

Chisholm NC, Packard AR, Koss WA, Juraska JM (2012) The effects of long-term treatment with estradiol and medroxyprogesterone acetate on tyrosine hydroxylase fibers and neuron number in the medial prefrontal cortex of aged female rats. *Endocrinology* 153:4874-4882.

Cunningham MG, Bhattacharyya S, Benes FM (2002) Amygdalo-cortical sprouting continues into early adulthood: implications for the development of normal and abnormal function during adolescence. *J Comp Neurol* 453:116-130.

Dalley JW, Cardinal RN, Robbins TW (2004) Prefrontal executive and cognitive functions in rodents: neural and neurochemical substrates. *Neurosci Biobehav Rev* 28:771-784.

- Descarries L, Lemay B, Doucet G, Berger B (1987) Regional and laminar density of the dopamine innervation in adult rat cerebral cortex. *Neuroscience* 21(3):807-24.
- Devoto P, Flore G (2006) On the origin of cortical dopamine: is it a co-transmitter in noradrenergic neurons? *Curr Neuroparmacol* 4(2):115-25.
- Dunlop BW, Nemeroff CB (2007) The role of dopamine in the pathophysiology of depression. *Arch Gen Psychiatry* 64(3):327-337.
- Emson PC, Koob GF (1978) The origin and distribution of dopamine-containing afferents to the rat frontal cortex. *Brain Res* 142:249-267.
- Forster B, Van De Ville D, Berent J, Sage D, Unser M (2004) Complex wavelets for extended depth-of-field: a new method for the fusion of multichannel microscopy images. *Microsc Res Tech* 65(1-2):33-42.
- Giedd JN, Blumenthal J, Jeffries NO, Castellanos FX, Liu H, Zijdenbos A, Paus T, Evans AC, Rapoport JL (1999) Brain development during childhood and adolescence: a longitudinal MRI study. *Nat Neurosci* 2(10):861-863.
- Giedd JN, Kechavan M, Paus T (2008) Why do many psychiatric disorders emerge during adolescence? *Nat Rev Neurosci* 9(12):947-957.
- Gillies GE, Virdee K, McArthur S, Dalley JW (2014) Sex-dependent diversity in ventral tegmental dopaminergic neurons and developmental programming: a molecular, cellular and behavioral analysis. *Neuroscience* 282:69-85.
- Gruber AJ, Calhoon GG, Shusterman I, Schoenbaum G, Roesch MR, O'Donnell P (2010) More is less: a disinhibited prefrontal cortex impairs cognitive flexibility. *J Neurosci* 30(50):17102-17110.
- Huttenlocher PR, Dabholkar AS (1997) Regional differences in synaptogenesis in human cerebral cortex. *J Comp Neurol* 387:167-178.
- Johnson ML, Day AE, Ho CC, Walker QD, Francis R, Kuhn CM (2010) Androgen decreases dopamine neurone survival in rat midbrain. *J Neuroendocrinol* 22(4):238-247.
- Juraska JM, Sisk CL, DonCarlos LL (2013) Sexual differentiation of the adolescent rodent brain: Hormonal influences and developmental mechanisms. *Horm Behav* 64:203-210.
- Kalsbeek A, Voorn P, Buijs RM, Pool CW, Uylings HBM (1988) Development of the dopaminergic innervation in the prefrontal cortex of the rat. *J Comp Neurol* 269:58-72.

Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE (2005) Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62(6):593-602.

Koss WA, Lloyd MM, Sadowski RN, Wise LM, Juraska JM (2015) Gonadectomy before puberty increases the number of neurons and glia in the medial prefrontal cortex of female, but not male, rats. *Dev Psychobiol* 57(3):305-12.

Kritzer MF, Brewer A, Montalman F, Davenport M, Robinson JK (2007) Effects of gonadectomy on performance in operant tasks measuring prefrontal cortical function in adult male rats. *Horm Behav* 51:183-194.

Kritzer MF, Creutz LM (2008) Region and sex differences in constituent dopamine neurons and immunoreactivity for intracellular estrogen and androgen receptors in mesocortical projections in rats. *J Neurosci* 28(38):9525-9535.

Levitt P, Moore RY (1979) Development of the noradrenergic innervation of neocortex. *Brain Res* 162:243-259.

Lewis DA (1997) Development of the prefrontal cortex during adolescence: Insights into vulnerable neural circuits in schizophrenia. *Neuropsychopharmacology* 16(6):385-398.

Lewis DA, Campbell MJ, Foote SL, Goldstein M, Morrison JH (1987) The distribution of tyrosine hydroxylase-immunoreactive fibers in primate neocortex is widespread but regionally specific. *J Neurosci* 7(1):279-290.

Markham JA, Morris JR, Juraska JM (2007) Neuron number decreases in the rat ventral, but not dorsal, medial prefrontal cortex between adolescence and adulthood. *Neuroscience* 144:961-968.

Miner LH, Schroeter S, Blakely RD, Sesack SR (2003) Ultrastructural localization of the norepinephrine transporter in superficial and deep layers of the rat prelimbic prefrontal cortex and its spatial relationship to probable dopamine terminals. *J Comp Neurol* 466:478-494.

Money KM, Stanwood GD (2013) Developmental origins of brain disorders: roles for dopamine. *Front Cell Neurosci* 7:260.

Monroe SM, Harkness KL (2005) Life stress, the "kindling" hypothesis, and the recurrence of depression: considerations from a life stress perspective. *Psychol Rev* 112(2):417-445.

Naneix F, Marchand AR, Di Scala G, Pape JR, Coutureau E (2012) Parallel maturation of goal-directed behavior and dopaminergic systems during adolescence. *J Neurosci* 32(46):16223-16232.

Palazidou E (2012) The neurobiology of depression. *Br Med Bull* 101:127-145.

Parker G, Brotchie H (2010) Gender differences in depression. *Int Rev Psychiatry* 22(5):429-436.

Satoh J, Suzuki K (1990) Tyrosine hydroxylase-immunoreactive neurons in the mouse cerebral cortex during the postnatal period. *Dev Brain Res* 53:1-5.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image analysis. *Nat Meth* 9(7):676-682.

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Meth* 9:671-675.

Spear LP (2000) The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev* 24:417-463.

Steinberg L (2010) A dual systems model of adolescent risk-taking. *Dev Psychobiol* 52:216-224.

Tseng KY, O'Donnell P (2007) Dopamine modulation of prefrontal cortical interneurons changes during adolescence. *Cereb Cortex* 17(5):1235-1240.

Verney C, Berger B, Adrien J, Vigny A, Gay M (1982) Development of the dopaminergic innervation of the rat cerebral cortex. A light microscopic immunocytochemical study using anti-tyrosine hydroxylase antibodies. *Dev Brain Res* 5:41-52.

Verney C, Milosevic A, Alvarez C, Berger B (1993) Immunocytochemical evidence of well-developed dopaminergic and noradrenergic innervations in the frontal cerebral cortex of human fetuses at midgestation. *J Comp Neurol* 336:331-334.

Zhang J, Darling RD, Paul IA, Simpson KL, Chen K, Shih JC, Lin RC (2011) Altered expression of tyrosine hydroxylase in the locus coeruleus noradrenergic system in citalopram neonatally exposed rats and monoamine oxidase a knock out mice. *Anat Rec* 294(10):1685-97.